

ONOFRI INEQUALITIES AND RIGIDITY RESULTS

JEAN DOLBEAULT* AND MARIA J. ESTEBAN

Ceremade, CNRS UMR n° 7534 and Université Paris-Dauphine
PSL research university
Place de Lattre de Tassigny, 75775 Paris Cédex 16, France

GASPARD JANKOWIAK

RICAM and Universität Wien
Wolfgang Pauli Institute
Oskar-Morgenstern-Platz 1, 1090 Wien, Austria

(Communicated by José A. Carrillo)

ABSTRACT. This paper is devoted to the Moser-Trudinger-Onofri inequality on smooth compact connected Riemannian manifolds. We establish a rigidity result for the Euler-Lagrange equation and deduce an estimate of the optimal constant in the inequality on two-dimensional closed Riemannian manifolds. Compared to existing results, we provide a non-local criterion which is well adapted to variational methods, introduce a nonlinear flow along which the evolution of a functional related with the inequality is monotone and get an integral remainder term which allows us to discuss optimality issues. As an important application of our method, we also consider the non-compact case of the Moser-Trudinger-Onofri inequality on the two-dimensional Euclidean space, with weights. The standard weight is the one that is computed when projecting the two-dimensional sphere using the stereographic projection, but we also give more general results which are of interest, for instance, for the Keller-Segel model in chemotaxis.

In this paper we assume that (\mathcal{M}, g) is a smooth compact connected Riemannian manifold of dimension $d \geq 1$, without boundary. We denote by Δ the Laplace-Beltrami operator on \mathcal{M} . For simplicity, we assume that the volume of \mathcal{M} , is chosen equal to 1 and use the notation dv_g for the volume element. We shall also denote by Ric the Ricci tensor, by $\text{H}u$ the Hessian of u and by

$$Lu := \text{H}u - \frac{g}{d} \Delta u$$

the trace free Hessian. Let us denote by $M[u]$ the trace free tensor

$$M[u] := \nabla u \otimes \nabla u - \frac{g}{d} |\nabla u|^2.$$

2010 *Mathematics Subject Classification*. Primary: 58J35, 58J05, 53C21; Secondary: 35J60, 35K55.

Key words and phrases. Moser-Trudinger-Onofri inequality, compact Riemannian manifold, Laplace-Beltrami operator, Ricci tensor, semilinear elliptic equation, rigidity, carré du champ, uniqueness, nonlinear diffusion, Sobolev inequality, Poincaré inequality, optimal constant.

* Corresponding author: Jean Dolbeault.

Let $g^{i,j}$ be the inverse of the metric tensor, *i.e.*, $g^{i,j}g_{j,k} = \delta_k^i$, where δ_k^i denotes the Kronecker symbol. If $A_{i,j}$ and $B_{i,j}$ are two tensors, we use the notation

$$A : B := g^{i,m}g^{j,n}A_{i,j}B_{m,n} \quad \text{and} \quad \|A\|^2 := A : A,$$

where we used the Einstein summation convention. We define

$$\lambda_* := \inf_{u \in H^2(\mathcal{M}) \setminus \{0\}} \frac{\int_{\mathcal{M}} \left[\|Lu - \frac{1}{2}M[u]\|^2 + \text{Ric}(\nabla u, \nabla u) \right] e^{-u/2} dv_g}{\int_{\mathcal{M}} |\nabla u|^2 e^{-u/2} dv_g}. \quad (1)$$

In this paper we will prove the two following results.

Theorem 1. *Assume that $d = 2$ and $\lambda_* > 0$. If u is a smooth solution to*

$$-\frac{1}{2}\Delta u + \lambda = e^u, \quad (2)$$

then u is a constant function if $\lambda \in (0, \lambda_)$.*

Note that the hypothesis that $\lambda_* > 0$ is in principle weaker than assuming that the Ricci curvature is everywhere positive on \mathcal{M} . See Remark 5 for more details.

Next, let us consider the Moser-Trudinger-Onofri inequality on \mathcal{M} written as

$$\frac{1}{4} \|\nabla u\|_{L^2(\mathcal{M})}^2 + \lambda \int_{\mathcal{M}} u dv_g \geq \lambda \log \left(\int_{\mathcal{M}} e^u dv_g \right) \quad \forall u \in H^1(\mathcal{M}), \quad (3)$$

for some constant $\lambda > 0$. Let us denote by λ_1 the first positive eigenvalue of $-\Delta$. We would like to draw the attention of the reader to the fact that, because of the normalization of the volume of \mathcal{M} , there is a discrepancy of a factor 4π with many results that are available in the literature. This factor corresponds to the surface of the usual \mathbb{S}^2 sphere, considered with the measure induced by Lebesgue's measure in \mathbb{R}^3 .

Corollary 2. *If $d = 2$, then (3) holds with $\lambda = \Lambda := \min\{1, \lambda_*\} \leq \lambda_1/2$. Moreover, if Λ is strictly smaller than $\lambda_1/2$, then the optimal constant in (3) is strictly larger than Λ .*

As we shall see later, in the case of the normalized sphere, $\lambda_* = \lambda_1/2 = 1$ is optimal, but for $\lambda = \lambda_*$, Eq. (2) has non-constant solutions because of the conformal invariance: see [31, 24] for more details on the Moser-Trudinger-Onofri inequality on the sphere, and references therein. The interested reader is invited to refer to the historical papers [48, 38, 41], and to [27, 28] for recent results on functionals related to the inequality, that have been obtained by variational methods. These last two papers solve the question, in any dimension, of whether the *first best constant* can be reached. This is equivalent to showing that the difference of the two terms in (3) is bounded from below. Earlier results have been obtained by T. Aubin in [1], in the case of the sphere \mathbb{S}^n , and P. Cherrier in [21] for general 2-manifolds. The present paper focuses on the value of the *second best constant*, defined as the largest value of λ such that (3) holds. The value of the *first best constant* is of little concern to us, as it appears as the $\frac{1}{4}$ coefficient in front of $\|\nabla u\|_{L^2(\mathcal{M})}^2$ and can be factored into λ . The method used by Z. Faget in [27, 28] relies on a blow-up analysis which is reminiscent for $d = 2$ of [40]. It generalizes some results contained in [2, Theorem 2.50 page 68]. Other references of general interest in the context of the Moser-Trudinger-Onofri inequality are [14, 5, 34, 42, 31]. A review of results related with the Moser-Trudinger-Onofri inequality in the case $\mathcal{M} = \mathbb{S}^2$

can be found in [24]. Let us mention that in [30], A. Ghigi provides a proof of this inequality based on the Prékopa-Leindler inequality and that many more details can be found in the book [31, Chapters 16-18] of N. Ghoussoub and A. Moradifam. In the context of Einstein-Kähler geometry another proof appears in [44, Theorem 5.2] (also see for instance [47, 43] and [8] for recent results on Kählerian manifolds). In [44], Y.A. Rubinstein gives a proof of the Onofri inequality on \mathbb{S}^2 that does not use symmetrization/rearrangement arguments. Also see [45] and in particular [45, Corollary 10.12] which contains a reinforced version of the inequality. We shall refer to [17] for background material in this direction. The reader interested in understanding how the Moser-Trudinger-Onofri inequality is related to the problem of prescribing the Gaussian curvature on \mathbb{S}^2 is invited to refer to [15, Section 3] for an introductory survey, and to [16, 18, 19] for more details. There are also many references about the equation (2), which is a particular version of the so-called Liouville equation. The book [46] contains a good description of the literature about this equation and many references. More references will be given within the text, whenever needed.

At this point, we should emphasize that in most of the literature the Moser-Trudinger-Onofri inequality in dimension $d = 2$ is not written as in (3), but in the form

$$e^{\mu_2 \|\nabla u\|_{L^2(\mathcal{M})}^2} \geq C \int_{\mathcal{M}} e^u dv_g$$

for all functions $u \in H^1(\mathcal{M})$ such that $\int_{\mathcal{M}} u dv_g = 0$, for some constant C which is in general non-explicit. In dimension $d = 2$, the optimal constant is $\mu_2 = \frac{1}{4}$. This amounts to write that the functional

$$u \mapsto \mu_2 \|\nabla u\|_{L^2(\mathcal{M})}^2 + \int_{\mathcal{M}} u dv_g - \log \left(\int_{\mathcal{M}} e^u dv_g \right)$$

is bounded from below by $\log C$. The issue of the *first best constant* is to prove that if μ_2 is replaced by a smaller constant, the functional becomes unbounded from below. This is different than proving Inequality (3), except when $C = 1$ and $\lambda = 1$. E. Onofri proved in [41] that this is the case, with optimal values for both C and λ , when $\mathcal{M} = \mathbb{S}^2$, up to a factor 4π that comes from the normalization of $\text{vol}_g(\mathcal{M})$. Except for the sphere we are aware of only one occurrence in the literature of the form (3) of the inequality, that has been derived by E. Fontenas in [29, Théorème 2] under more restrictive conditions on \mathcal{M} . This result will be commented in more detail in Remark 4.

The proof of Theorem 1 is a *rigidity* method inspired by the one of [9] for the equation

$$-\Delta u + \lambda u = u^{p-1},$$

which is the Euler-Lagrange equation corresponding to the optimality case in the interpolation inequality

$$\|\nabla v\|_{L^2(\mathcal{M})}^2 \geq \frac{\lambda}{p-2} \left[\|v\|_{L^p(\mathcal{M})}^2 - \|v\|_{L^2(\mathcal{M})}^2 \right] \quad \forall v \in H^1(\mathcal{M}). \tag{4}$$

See [9, 37, 3, 22, 25] for further results on this problem and [2, 33] for general accounts on Sobolev’s inequality on Riemannian manifolds. Concerning spectral issues, a standard textbook is [7].

The case of the exponential nonlinearity in (2) has been much less considered in the literature, except when \mathcal{M} is the two dimensional sphere \mathbb{S}^2 . Let us mention the uniqueness result of [20] for (2) with $\lambda = 1$. In [29], and in [6] in the case of

the ultraspherical operator, the result is achieved by considering the interpolation inequalities (4) and then, as in [9] or [5] (in the case of the sphere), by taking the limit as $p \rightarrow \infty$. Here we consider a direct approach, based on rigidity methods and an associated nonlinear flow. As far as we know, this is an entirely new approach which is interesting in that it provides explicit estimates on the optimal constant in (3).

One may wonder if rigidity results can be achieved for dimensions $d > 2$ with our method. We will give a negative answer in Section 1. Corollary 2 is established in Section 2 using a nonlinear flow that has already been considered on the sphere in [24]. The case $d = 1$ is very simple and will be considered for the sake of completeness in Section 3. An important application of our method is the case of the Euclidean space with weights, with applications to chemotaxis. Section 4 is devoted to this issue with a main result in this direction stated in Theorem 8, that raises difficult questions of symmetry breaking.

1. Proof of Theorem 1. In this section we consider a smooth solution to (2) and perform a computation to prove the rigidity result of Theorem 1. There is no *a priori* reason to assume that $d = 2$ and so we shall do the computations for any dimension $d \geq 1$, which raises no special additional difficulties. However, due to restrictions that are inherent to the method and will be explicitly exposed, only $d = 2$ can be covered. On several occasions, one has to divide by $(d - 1)$, so the case $d = 1$ has to be excluded and will be handled directly in Section 3.

In the case of (4), it is well known (see [37, 3, 26]) that an interpolation depending on a parameter $\theta \in (0, 1)$ between an estimate based on the Ricci curvature and another one based on the first eigenvalue of the Laplace-Beltrami operator can be used to obtain some improvements. Here we apply the same technique and realize in the end that only $\theta = 1$ is admissible in dimension $d = 2$. However, when d is considered as a real parameter in the range $(1, 2)$, it is possible to optimize on θ when $0 \leq \theta \leq 1$. We will comment this and possible improvements at the end of this section.

Preliminaries. A simple expansion of the square shows that

$$\|Hu\|^2 = \|Lu\|^2 + \frac{1}{d} (\Delta u)^2.$$

The Bochner-Lichnerovitz-Weitzenböck formula asserts that

$$\frac{1}{2} \Delta |\nabla u|^2 = \|Hu\|^2 + \nabla(\Delta u) \cdot \nabla u + \text{Ric}(\nabla u, \nabla u)$$

where Ric denotes the Ricci tensor and, as a consequence,

$$\frac{1}{2} \Delta |\nabla u|^2 = \|Lu\|^2 + \frac{1}{d} (\Delta u)^2 + \nabla(\Delta u) \cdot \nabla u + \text{Ric}(\nabla u, \nabla u). \quad (5)$$

An identity based on integrations by parts. Using integrations by parts, we may notice that

$$\begin{aligned} & - \int_{\mathcal{M}} \Delta u |\nabla u|^2 e^{-u/2} dv_g \\ &= -\frac{1}{2} \int_{\mathcal{M}} |\nabla u|^4 e^{-u/2} dv_g + 2 \int_{\mathcal{M}} Hu : \nabla u \otimes \nabla u e^{-u/2} dv_g \end{aligned}$$

$$\begin{aligned}
 &= -\frac{1}{2} \int_{\mathcal{M}} |\nabla u|^4 e^{-u/2} dv_g + 2 \int_{\mathcal{M}} \left(Lu + \frac{g}{d} \Delta u \right) : \nabla u \otimes \nabla u e^{-u/2} dv_g \\
 &= -\frac{1}{2} \int_{\mathcal{M}} |\nabla u|^4 e^{-u/2} dv_g + 2 \int_{\mathcal{M}} Lu : M[u] e^{-u/2} dv_g \\
 &\quad + \frac{2}{d} \int_{\mathcal{M}} \Delta u |\nabla u|^2 e^{-u/2} dv_g,
 \end{aligned}$$

with $M[u] := \nabla u \otimes \nabla u - \frac{g}{d} |\nabla u|^2$, which proves that

$$\begin{aligned}
 \frac{d+2}{d} \int_{\mathcal{M}} \Delta u |\nabla u|^2 e^{-u/2} dv_g \\
 = \frac{1}{2} \int_{\mathcal{M}} |\nabla u|^4 e^{-u/2} dv_g - 2 \int_{\mathcal{M}} Lu : M[u] e^{-u/2} dv_g
 \end{aligned}$$

and finally

$$\begin{aligned}
 \int_{\mathcal{M}} \Delta u |\nabla u|^2 e^{-u/2} dv_g \\
 = \frac{1}{2} \frac{d}{d+2} \int_{\mathcal{M}} |\nabla u|^4 e^{-u/2} dv_g - \frac{2d}{d+2} \int_{\mathcal{M}} Lu : M[u] e^{-u/2} dv_g. \quad (6)
 \end{aligned}$$

An identity based on the Bochner-Lichnerovicz-Weitzenböck formula. By expanding $\Delta(e^{-u/2}) = (\frac{1}{4} |\nabla u|^2 - \frac{1}{2} \Delta u) e^{-u/2}$, we have that

$$\int_{\mathcal{M}} |\nabla u|^2 \Delta(e^{-u/2}) dv_g = \frac{1}{4} \int_{\mathcal{M}} |\nabla u|^4 e^{-u/2} dv_g - \frac{1}{2} \int_{\mathcal{M}} \Delta u |\nabla u|^2 e^{-u/2} dv_g$$

so that, if we multiply (5) by $e^{-u/2}$ and integrate by parts, then we get

$$\begin{aligned}
 &\frac{1}{8} \int_{\mathcal{M}} |\nabla u|^4 e^{-u/2} dv_g - \frac{1}{4} \int_{\mathcal{M}} \Delta u |\nabla u|^2 e^{-u/2} dv_g \\
 &= \int_{\mathcal{M}} \|Lu\|^2 e^{-u/2} dv_g + \frac{1}{d} \int_{\mathcal{M}} (\Delta u)^2 e^{-u/2} dv_g \\
 &\quad + \int_{\mathcal{M}} (\nabla(\Delta u) \cdot \nabla u) e^{-u/2} dv_g + \int_{\mathcal{M}} Ric(\nabla u, \nabla u) e^{-u/2} dv_g \\
 &= \int_{\mathcal{M}} \|Lu\|^2 e^{-u/2} dv_g - \frac{d-1}{d} \int_{\mathcal{M}} (\Delta u)^2 e^{-u/2} dv_g \\
 &\quad + \frac{1}{2} \int_{\mathcal{M}} \Delta u |\nabla u|^2 e^{-u/2} dv_g + \int_{\mathcal{M}} Ric(\nabla u, \nabla u) e^{-u/2} dv_g,
 \end{aligned}$$

from which we deduce that

$$\begin{aligned}
 &\int_{\mathcal{M}} (\Delta u)^2 e^{-u/2} dv_g \\
 &= \frac{3}{4} \frac{d}{d-1} \int_{\mathcal{M}} \Delta u |\nabla u|^2 e^{-u/2} dv_g - \frac{1}{8} \frac{d}{d-1} \int_{\mathcal{M}} |\nabla u|^4 e^{-u/2} dv_g \\
 &\quad + \frac{d}{d-1} \int_{\mathcal{M}} \|Lu\|^2 e^{-u/2} dv_g + \frac{d}{d-1} \int_{\mathcal{M}} Ric(\nabla u, \nabla u) e^{-u/2} dv_g. \quad (7)
 \end{aligned}$$

A Poincaré inequality. Since

$$4\Delta(e^{-u/4}) = \frac{1}{4}|\nabla u|^2 e^{-u/4} - \Delta u e^{-u/4},$$

we get that

$$\begin{aligned} 16 \int_{\mathcal{M}} |\Delta(e^{-u/4})|^2 dv_g &= \frac{1}{16} \int_{\mathcal{M}} |\nabla u|^4 e^{-u/2} dv_g - \frac{1}{2} \int_{\mathcal{M}} |\nabla u|^2 \Delta u e^{-u/2} dv_g \\ &\quad + \int_{\mathcal{M}} (\Delta u)^2 e^{-u/2} dv_g. \end{aligned}$$

On the other hand, a Poincaré inequality applied to $\nabla(e^{-u/4})$, as in [26, Lemma 7], shows that

$$\int_{\mathcal{M}} |\Delta(e^{-u/4})|^2 dv_g \geq \lambda_1 \int_{\mathcal{M}} |\nabla(e^{-u/4})|^2 dv_g = \frac{\lambda_1}{16} \int_{\mathcal{M}} |\nabla u|^2 e^{-u/2} dv_g, \quad (8)$$

so that

$$\begin{aligned} \int_{\mathcal{M}} (\Delta u)^2 e^{-u/2} dv_g &\geq \lambda_1 \int_{\mathcal{M}} |\nabla u|^2 e^{-u/2} dv_g - \frac{1}{16} \int_{\mathcal{M}} |\nabla u|^4 e^{-u/2} dv_g \\ &\quad + \frac{1}{2} \int_{\mathcal{M}} |\nabla u|^2 \Delta u e^{-u/2} dv_g. \quad (9) \end{aligned}$$

An identity based on the equation. By expanding $\Delta(e^{-u/2}) = (\frac{1}{4}|\nabla u|^2 - \frac{1}{2}\Delta u)e^{-u/2}$, we have that

$$\int_{\mathcal{M}} (-\frac{1}{2}\Delta u)\Delta(e^{-u/2}) dv_g = \frac{1}{4} \int_{\mathcal{M}} (\Delta u)^2 e^{-u/2} dv_g - \frac{1}{8} \int_{\mathcal{M}} \Delta u |\nabla u|^2 e^{-u/2} dv_g$$

so that, if we multiply (2) by $\Delta(e^{-u/2}) - \frac{1}{2}|\nabla u|^2 e^{-u/2}$, then we get

$$\frac{1}{4} \int_{\mathcal{M}} (\Delta u)^2 e^{-u/2} dv_g + \frac{1}{8} \int_{\mathcal{M}} \Delta u |\nabla u|^2 e^{-u/2} dv_g - \frac{\lambda}{2} \int_{\mathcal{M}} |\nabla u|^2 e^{-u/2} dv_g = 0.$$

We then split the first term in the equality as the sum of $\frac{1-\theta}{4} \int_{\mathcal{M}} (\Delta u)^2 e^{-u/2} dv_g$ and $\frac{\theta}{4} \int_{\mathcal{M}} (\Delta u)^2 e^{-u/2} dv_g$, with a parameter $\theta \leq 1$, and we bound the resulting terms using (9) and (7), respectively:

$$\begin{aligned} &\frac{1-\theta}{4} \left[\lambda_1 \int_{\mathcal{M}} |\nabla u|^2 e^{-u/2} dv_g - \frac{1}{16} \int_{\mathcal{M}} |\nabla u|^4 e^{-u/2} dv_g \right. \\ &\quad \left. + \frac{1}{2} \int_{\mathcal{M}} |\nabla u|^2 \Delta u e^{-u/2} dv_g \right] \\ &+ \frac{\theta}{4} \left[\frac{3}{4} \frac{d}{d-1} \int_{\mathcal{M}} \Delta u |\nabla u|^2 e^{-u/2} dv_g - \frac{1}{8} \frac{d}{d-1} \int_{\mathcal{M}} |\nabla u|^4 e^{-u/2} dv_g \right. \\ &\quad \left. + \frac{d}{d-1} \int_{\mathcal{M}} \|Lu\|^2 e^{-u/2} dv_g + \frac{d}{d-1} \int_{\mathcal{M}} \text{Ric}(\nabla u, \nabla u) e^{-u/2} dv_g \right] \\ &\quad + \frac{1}{8} \int_{\mathcal{M}} \Delta u |\nabla u|^2 e^{-u/2} dv_g - \frac{\lambda}{2} \int_{\mathcal{M}} |\nabla u|^2 e^{-u/2} dv_g \leq 0. \end{aligned}$$

Collecting terms, we get

$$\begin{aligned} & \frac{\theta}{4} \frac{d}{d-1} \left[\int_{\mathcal{M}} \|Lu\|^2 e^{-u/2} dv_g + \int_{\mathcal{M}} \text{Ric}(\nabla u, \nabla u) e^{-u/2} dv_g \right] \\ & - \frac{1}{64} \left(1 - \theta + 2\theta \frac{d}{d-1} \right) \int_{\mathcal{M}} |\nabla u|^4 e^{-u/2} dv_g \\ & + \left(\frac{1-\theta}{8} + \frac{3\theta}{16} \frac{d}{d-1} + \frac{1}{8} \right) \int_{\mathcal{M}} \Delta u |\nabla u|^2 e^{-u/2} dv_g \\ & + \left(\frac{1-\theta}{4} \lambda_1 - \frac{\lambda}{2} \right) \int_{\mathcal{M}} |\nabla u|^2 e^{-u/2} dv_g \leq 0 \end{aligned}$$

and can now use (6) to obtain

$$\begin{aligned} & \frac{\theta}{4} \frac{d}{d-1} \left[\int_{\mathcal{M}} \|Lu\|^2 e^{-u/2} dv_g + \int_{\mathcal{M}} \text{Ric}(\nabla u, \nabla u) e^{-u/2} dv_g \right] \\ & - \frac{1}{64} \left(1 - \theta + 2\theta \frac{d}{d-1} \right) \int_{\mathcal{M}} |\nabla u|^4 e^{-u/2} dv_g \\ & + \left(\frac{1-\theta}{8} + \frac{3\theta}{16} \frac{d}{d-1} + \frac{1}{8} \right) \left[\frac{1}{2} \frac{d}{d+2} \int_{\mathcal{M}} |\nabla u|^4 e^{-u/2} dv_g \right. \\ & \quad \left. - \frac{2d}{d+2} \int_{\mathcal{M}} Lu : M[u] e^{-u/2} dv_g \right] \\ & + \left(\frac{1-\theta}{4} \lambda_1 - \frac{\lambda}{2} \right) \int_{\mathcal{M}} |\nabla u|^2 e^{-u/2} dv_g \leq 0. \end{aligned}$$

Recall that $M[u]$ denotes the trace free tensor

$$M[u] := \nabla u \otimes \nabla u - \frac{g}{d} |\nabla u|^2.$$

We observe that

$$\|M[u]\|^2 = \left\| \nabla u \otimes \nabla u - \frac{g}{d} |\nabla u|^2 \right\|^2 = \left(1 - \frac{1}{d} \right) |\nabla u|^4.$$

Altogether we end up with

$$\begin{aligned} & \int_{\mathcal{M}} \left(\mathbf{a} \|Lu\|^2 + \mathbf{b} (Lu : M[u]) + \mathbf{c} \|M[u]\|^2 \right) e^{-u/2} dv_g \\ & + \frac{\theta}{4} \frac{d}{d-1} \int_{\mathcal{M}} \text{Ric}(\nabla u, \nabla u) e^{-u/2} dv_g \\ & + \left(\frac{1-\theta}{4} \lambda_1 - \frac{\lambda}{2} \right) \int_{\mathcal{M}} |\nabla u|^2 e^{-u/2} dv_g \leq 0 \end{aligned}$$

with

$$\begin{aligned} \mathbf{a} &= \frac{\theta}{4} \frac{d}{d-1}, \\ \mathbf{b} &= - \left(\frac{1-\theta}{8} + \frac{3\theta}{16} \frac{d}{d-1} + \frac{1}{8} \right) \frac{2d}{d+2}, \\ \mathbf{c} &= \left[\left(\frac{1-\theta}{8} + \frac{3\theta}{16} \frac{d}{d-1} + \frac{1}{8} \right) \frac{1}{2} \frac{d}{d+2} - \frac{1}{64} \left(1 - \theta + 2\theta \frac{d}{d-1} \right) \right] \frac{d}{d-1}. \end{aligned}$$

Remark 3. By the Lichnerowicz' theorem (see [36] or [26, Section 2])

$$\frac{d}{d-1} \int_{\mathcal{M}} \operatorname{Ric}(\nabla u, \nabla u) e^{-u/2} dv_g \leq \int_{\mathcal{M}} |\nabla u|^2 e^{-u/2} dv_g$$

so that the largest possible value of λ for which we *a priori* know that

$$\frac{\theta}{4} \frac{d}{d-1} \int_{\mathcal{M}} \operatorname{Ric}(\nabla u, \nabla u) e^{-u/2} dv_g + \left(\frac{1}{4} \lambda_1 (1-\theta) - \lambda\right) \int_{\mathcal{M}} |\nabla u|^2 e^{-u/2} dv_g$$

is nonnegative corresponds to the smallest possible value of θ , *i.e.* $\theta = \theta_0(d)$.

Discussion. With a simple but tedious computation, one can show that the discriminant $\delta := b^2 - 4ac$ has the sign of

$$16(d-1)^2 - (6-d)(d+2)\theta.$$

If we denote by $\theta_0 = \theta_0(d)$ the value of θ for which $\delta = 0$, then we have

$$\theta_0 = \frac{16(d-1)^2}{(6-d)(d+2)}.$$

Altogether we can rewrite our estimate as

$$\begin{aligned} & \mathbf{a} \int_{\mathcal{M}} \left\| Lu + \frac{\mathbf{b}}{2\mathbf{a}} M[u] \right\|^2 e^{-u/2} dv_g + \left(\mathbf{c} - \frac{\mathbf{b}^2}{4\mathbf{a}} \right) \int_{\mathcal{M}} \|M[u]\|^2 e^{-u/2} dv_g \\ & + 4(1-\theta) \int_{\mathcal{M}} |\Delta(e^{-u/4})|^2 dv_g + \frac{\theta}{4} \frac{d}{d-1} \int_{\mathcal{M}} \operatorname{Ric}(\nabla u, \nabla u) e^{-u/2} dv_g \\ & - \frac{\lambda}{2} \int_{\mathcal{M}} |\nabla u|^2 e^{-u/2} dv_g = 0 \end{aligned}$$

and use the Poincaré inequality (8) to establish that

$$\begin{aligned} 0 & \geq \mathbf{a} \int_{\mathcal{M}} \left\| Lu + \frac{\mathbf{b}}{2\mathbf{a}} M[u] \right\|^2 e^{-u/2} dv_g - \frac{\delta}{4\mathbf{a}} \int_{\mathcal{M}} \|M[u]\|^2 e^{-u/2} dv_g \\ & + \left(\frac{1}{4} \lambda_1 (1-\theta) - \frac{\lambda}{2}\right) \int_{\mathcal{M}} |\nabla u|^2 e^{-u/2} dv_g + \frac{\theta}{4} \frac{d}{d-1} \int_{\mathcal{M}} \operatorname{Ric}(\nabla u, \nabla u) e^{-u/2} dv_g. \end{aligned} \tag{10}$$

Our goal is to show that u has to be a constant, that is, $\int_{\mathcal{M}} |\nabla u|^2 e^{-u/2} dv_g = 0$. We assume that $d \geq 2$ is an integer. The discriminant δ is nonpositive if and only if $d < 6$ and $\theta \geq \theta_0(d)$. This is compatible with the condition $\theta \leq 1$ only if $d = 2$; in that case, $\theta = \theta_0(2) = 1$ and as a consequence, we can rewrite (10) as

$$\begin{aligned} 0 & \geq \int_{\mathcal{M}} \left\| Lu - \frac{1}{2} M[u] \right\|^2 e^{-u/2} dv_g + \int_{\mathcal{M}} \operatorname{Ric}(\nabla u, \nabla u) e^{-u/2} dv_g \\ & - \lambda \int_{\mathcal{M}} |\nabla u|^2 e^{-u/2} dv_g \\ & \geq (\lambda_* - \lambda) \int_{\mathcal{M}} |\nabla u|^2 e^{-u/2} dv_g. \end{aligned}$$

Hence we have shown that $\nabla u \equiv 0$ for any $\lambda < \lambda_*$, which concludes the proof of Theorem 1.

Remark 4. In order to compare our results with the results deduced from the *curvature-dimension* method (see for instance [4]), we can consider the case where Ric is uniformly bounded from below by some positive constant ρ and formally assume that $d \in (1, 2)$ takes real values. This can be made precise for instance

in the setting of the ultra-spherical operator (see for instance [6]), with exactly the same conditions as above. See [29] and [24, Section 7.1] for more details. If $1 < d \leq 2$, we find that rigidity holds if

$$\lambda \leq \max_{\theta \in [\theta_0(d), 1]} \left(\frac{1}{2} \lambda_1 (1 - \theta) + \frac{\theta}{2} \frac{d}{d-1} \rho \right) = \frac{1}{2} \lambda_1 (1 - \theta_0(d)) + \frac{\theta_0(d)}{2} \frac{d}{d-1} \rho$$

according to Remark 3. Let $x = \frac{d}{d-1} \frac{\rho}{\lambda_1} \leq 1$. We have found that that rigidity holds if

$$2 \frac{\lambda}{\lambda_1} \leq 1 - \theta_0(d) + \theta_0(d) x =: f_1(x)$$

Quite surprisingly, a better condition has been obtained in [29, Théorème 2], when $1 < d < 2$, which amounts to

$$2 \frac{\lambda}{\lambda_1} \leq d(2 - d) + (d - 1)^2 x =: f_2(x),$$

by taking the limit as $p \rightarrow \infty$ in (4). We may indeed check that $f_2(x) - f_1(x) = \frac{(d-1)^2 (d-2)^2}{(6-d)(d+2)} (1 - x) \geq 0$

Without assuming the positivity of ρ , one gets a similar result with our approach. In the range $d \in (1, 2)$, our computations show that rigidity holds for any λ at most equal to the infimum on $u \in H^2(\mathcal{M}) \setminus \{0\}$ of

$$2 \int_{\mathcal{M}} \left[a \left\| Lu + \frac{b}{2a} M[u] \right\|^2 + 4(1-\theta) |\Delta(e^{-u/4})|^2 + \frac{\theta}{4} \frac{d}{d-1} \text{Ric}(\nabla u, \nabla u) \right] e^{-u/2} dv_g$$

under the condition that $\int_{\mathcal{M}} |\nabla u|^2 e^{-u/2} dv_g = 1$, $a = \frac{4d(d-1)}{(6-d)(d+2)}$, $b = -\frac{d(3d+2)}{2(6-d)(d+2)}$ and $\theta = \theta_0(d)$. However, in the same spirit as above, a passage to the limit as $p \rightarrow \infty$ in the inequality obtained in [26, Theorem 4] gives a better result.

Let us emphasize that these considerations are essentially formal because d is restricted to the interval $(1, 2)$ but can be entirely justified, as it is currently done in the *curvature-dimension* approach. See for instance [29, Théorème 2], and related references.

2. Proof of Corollary 2. Let us minimize the functional

$$\mathcal{F}_\lambda[u] := \frac{1}{4} \|\nabla u\|_{L^2(\mathcal{M})}^2 + \lambda \int_{\mathcal{M}} u dv_g - \lambda \log \left(\int_{\mathcal{M}} e^u dv_g \right)$$

on $H^1(\mathcal{M})$. According to [27, 28], $\mathcal{F}_\lambda[u]$ has no minimizer if $\lambda > 1$. Let us assume that

$$\lambda < 1$$

(we shall take care of the equality case later). It is then standard that there is a well-defined minimizer u . Note that since $\mathcal{F}_\lambda[u]$ does not change when adding some constant to u , we can choose the value of $\int_{\mathcal{M}} e^u dv_g = \lambda$ and then u satisfies (2). If u is smooth and $\lambda < \lambda_*$, we can apply the result of Theorem 1. Then the minimizer u has to be a constant, for instance $u \equiv 1$, so that

$$\mathcal{F}_\lambda[u] \geq \mathcal{F}_\lambda[1] = 0 \quad \forall u \in H^1(\mathcal{M}).$$

Notice that we can get rid of any smoothness requirement by considering the flow below. By passing to the limit as $\lambda \nearrow \lambda_*$, we get that the inequality also holds true if $\lambda = \lambda_*$.

Using the following Taylor expansion of \mathcal{F}_λ as $\varepsilon \rightarrow 0$,

$$\begin{aligned}\mathcal{F}_\lambda[1 + \varepsilon\varphi] &= \varepsilon^2 \left[\frac{1}{4} \|\nabla\varphi\|_{L^2(\mathcal{M})}^2 + \frac{\lambda}{2} \int_{\mathcal{M}} \varphi^2 dv_g \right] + o(\varepsilon^2) \\ &= \frac{\varepsilon^2}{2} \left(\frac{\lambda_1}{2} - \lambda \right) + o(\varepsilon^2),\end{aligned}$$

where φ is an eigenfunction associated with the first positive eigenvalue λ_1 of $-\Delta$, it is straightforward to see that the best constant λ in (3) is such that

$$\lambda \leq \frac{\lambda_1}{2}.$$

To complete the proof of Corollary 2, it remains to consider the case $\lambda_* < \lambda_1/2$ and show that the optimal constant cannot be equal to λ_* . This can be done in the same spirit as in [26, Corollary 2]. Let us consider the evolution equation defined by

$$\frac{\partial f}{\partial t} = \Delta(e^{-f/2}) - \frac{1}{2} |\nabla f|^2 e^{-f/2}, \quad (11)$$

with initial datum $u \in H^1(\mathcal{M})$. Let us define

$$\begin{aligned}\mathcal{G}_\lambda[f] := & \int_{\mathcal{M}} \|\mathbf{L}f - \frac{1}{2} \mathbf{M}[f]\|^2 e^{-f/2} dv_g + \int_{\mathcal{M}} \text{Ric}(\nabla f, \nabla f) e^{-f/2} dv_g \\ & - \lambda \int_{\mathcal{M}} |\nabla f|^2 e^{-f/2} dv_g.\end{aligned}$$

Then for any $\lambda \leq \lambda_*$ we have

$$\frac{d}{dt} \mathcal{F}_\lambda[f(t, \cdot)] = \int_{\mathcal{M}} \left(-\frac{1}{2} \Delta f + \lambda \right) \left(\Delta(e^{-f/2}) - \frac{1}{2} |\nabla f|^2 e^{-f/2} \right) dv_g = -\mathcal{G}_\lambda[f(t, \cdot)]$$

Since \mathcal{F}_λ is nonnegative and $\lim_{t \rightarrow \infty} \mathcal{F}_\lambda[f(t, \cdot)] = 0$, we obtain that

$$\mathcal{F}_\lambda[u] \geq \int_0^\infty \mathcal{G}_\lambda[f(t, \cdot)] dt$$

for any solution f to (11) with initial datum $u \in L^1(\mathcal{M})$ is such that $\nabla u \in L^2(\mathcal{M})$. We have an equality if the solution is smooth for any $t \geq 0$. Otherwise we have to regularize and then pass to the limit so that, with full generality, we can only expect an inequality.

Remark 5. One has to mention that the sphere $\mathcal{M} = \mathbb{S}^2$ is an important case of application of our method, for which other types of remainder terms can be produced. See [24] for more details. It has to be noted that on \mathbb{S}^2 we have $\lambda_* = \rho = \lambda_1/2 = 1$. As noted by many authors, *e.g.*, in [5, 6, 29, 24] (also see references in [24]), the Onofri inequality is a limit case of various Sobolev type inequalities, for which similar methods have been developed: see [25] for a review and some recent results.

Another interesting case for our method is the flat square torus, defined as the square $(0, 1) \times (0, 1)$ with double periodicity, or simply $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$. For general properties of the solutions of (2) on \mathbb{T}^2 , see [46, Section 2.5]. In this case, and also for any compact manifold \mathcal{M} with nonnegative Ricci curvature, by defining $v = e^{-u/2}$ one can see that

$$\lambda_* = \inf_{v \in H^2(\mathcal{M}) \setminus \{0\}} \frac{\int_{\mathcal{M}} \left(\frac{\|\mathbf{L}v\|^2}{v} + \frac{1}{v} \text{Ric}(\nabla v, \nabla v) \right) dv_g}{\int_{\mathcal{M}} \frac{|\nabla v|^2}{v} dv_g}.$$

According to [35, Theorem 0.1], there exists $C > 0$ such that for any $0 < \lambda < C$, all solutions u of (2) are uniformly bounded in $L^\infty(\mathcal{M})$. In the proofs of Theorem 1 and Corollary 2, it is henceforth possible to replace λ_\star by the infimum taken over the space of all functions $v \in H^2(\mathcal{M}) \setminus \{0\}$ such that $-2 \operatorname{var}(\log v) = \operatorname{var}(u) = \operatorname{supess}(u) - \operatorname{infess}(u) \leq \mathcal{K}$ for some positive constant \mathcal{K} , which is independent of $\lambda > 0$. Then, with this new definition of λ_\star , we obtain the estimate

$$\lambda_\star \geq e^{-\kappa/2} \inf \frac{\int_{\mathcal{M}} (\|Lv\|^2 + \operatorname{Ric}(\nabla v, \nabla v)) dv_g}{\int_{\mathcal{M}} |\nabla v|^2 dv_g} = e^{-\kappa/2} \inf \frac{\int_{\mathcal{M}} (\Delta v)^2 dv_g}{\int_{\mathcal{M}} |\nabla v|^2 dv_g},$$

where the last equality is a straightforward consequence of the Bochner-Lichnerowicz-Weitzenböck formula. As a consequence of the Poincaré inequality (see for instance [26, Lemma 5]), we obtain the estimate $\lambda_\star \geq e^{-\kappa/2} \lambda_1$. It is remarkable that our method applies when the lowest eigenvalue of $\operatorname{Ric}(\nabla v, \nabla v)$ takes value 0 in \mathcal{M} . In the case of the flat torus, we even have that $\operatorname{Ric}(\nabla v, \nabla v) \equiv 0$ on \mathcal{M} .

3. The case $d = 1$. For simplicity, we can consider the case of the circle. Hence we identify \mathcal{M} with the 1-periodic interval $[0, 1) \approx \mathbb{R}/\mathbb{Z} \approx \mathbb{S}^1$. Consider a solution of the ordinary differential equation

$$-\frac{1}{2} u'' + \lambda - e^u = 0 \tag{12}$$

with periodic boundary conditions. If we multiply the equation by $(e^{-u/2})'' - \frac{1}{2} |u'|^2 e^{-u/2}$, then we get that

$$\int_0^1 \left(\frac{1}{4} |u''|^2 + \frac{1}{8} |u'|^2 u'' - \frac{\lambda}{2} |u'|^2 \right) e^{-u/2} dx = 0.$$

The middle term is easy to handle using one integration by parts:

$$\int_0^1 |u'|^2 u'' e^{-u/2} dx = \frac{1}{6} \int_0^1 |u'|^4 e^{-u/2} dx. \tag{13}$$

Hence we have

$$\int_0^1 \left(\frac{1}{4} |u''|^2 + \frac{1}{48} |u'|^4 - \frac{\lambda}{2} |u'|^2 \right) e^{-u/2} dx = 0. \tag{14}$$

On the other hand, by the Poincaré inequality,

$$\int_0^1 \left| \left(e^{-u/4} \right)'' \right|^2 dx \geq 4\pi^2 \int_0^1 \left| \left(e^{-u/4} \right)' \right|^2 dx,$$

where $4\pi^2$ is the first positive eigenvalue of $-\frac{d^2}{dx^2}$ on the periodic interval of length 1. From (13) we derive

$$\int_0^1 \left(|u''|^2 - \frac{1}{48} |u'|^4 \right) e^{-u/2} dx - 4\pi^2 \int_0^1 |u'|^2 e^{-u/2} dx \geq 0. \tag{15}$$

Combining (14) and (15), we get

$$\frac{5}{96} \int_0^1 |u'|^4 e^{-u/2} dx + (2\pi^2 - \lambda) \int_0^1 |u'|^2 e^{-u/2} dx \leq 0.$$

Hence we have proven the following result.

Proposition 6. *Assume that $d = 1$. With the above notations, if u is a smooth solution to (12) on $\mathbb{S}^1 \approx [0, 1)$, then u is a constant function for any $\lambda \in (0, 2\pi^2)$.*

Exactly as in the case of a manifold of dimension two, a variational approach allows to deduce a Moser-Trudinger-Onofri inequality.

Corollary 7. *If $d = 1$, then the following inequality holds on $\mathbb{S}^1 \approx [0, 1)$:*

$$\frac{1}{8\pi^2} \int_0^1 |u'|^2 dx + \int_0^1 u dx \geq \log \left(\int_0^1 e^u dx \right) \quad \forall u \in H^1(\mathbb{S}^1).$$

Moreover $8\pi^2$ is the optimal constant.

The only difference with Corollary 2 is that we can identify the optimal constant in the inequality by considering $u = 1 + \varepsilon \varphi$ and by taking the limit as $\varepsilon \rightarrow 0$, with $\varphi(x) = \cos(2\pi x)$.

4. Weighted Moser-Trudinger-Onofri inequalities on the two-dimensional Euclidean space. The Euclidean Onofri inequality (see [14, 24]) can be deduced from (3) when $\mathcal{M} = \mathbb{S}^2$ using the stereographic projection and reads

$$\frac{1}{16\pi} \int_{\mathbb{R}^2} |\nabla u|^2 dx \geq \log \left(\int_{\mathbb{R}^2} e^u d\mu \right) - \int_{\mathbb{R}^2} u d\mu. \tag{16}$$

Here $d\mu = \mu(x) dx$ denotes the probability measure defined by $\mu(x) = \frac{1}{\pi} (1 + |x|^2)^{-2}$, $x \in \mathbb{R}^2$, and the inequality holds for any function $u \in L^1(\mathbb{R}^2, d\mu)$ such that $\nabla u \in L^2(\mathbb{R}^2)$. The constant 16π is optimal as can be shown by considering the inequality on \mathbb{S}^2 and comparing with the value given when expanding around a constant, as was done in Section 2.

In this section, our goal is to give sufficient conditions on a general probability density μ so that the inequality

$$\frac{1}{16\pi} \int_{\mathbb{R}^2} |\nabla u|^2 dx \geq \lambda \left[\log \left(\int_{\mathbb{R}^2} e^u d\mu \right) - \int_{\mathbb{R}^2} u d\mu \right] \tag{17}$$

holds for some $\lambda > 0$ and get an estimate of the optimal value of λ . Here $d\mu = \mu dx$ is a probability measure with density μ with respect to the Lebesgue measure. All our computations are done without symmetry assumption, and our final estimate is (30). In practical applications (see Examples 1–4) the function μ is radially symmetric *and* one has to assume that λ is in a range for which the solution to (17), or at least the optimal function for (16), is radially symmetric. This delicate issue of *symmetry breaking* will be illustrated in Example 3.

Since (17) does not change when adding some constant to u , we can look for minimizers satisfying the constraint $\int_{\mathbb{R}^2} e^u d\mu = 1$. These solve the Euler-Lagrange equation

$$-\frac{1}{8\pi} \Delta u + \lambda \mu - \lambda e^u \mu = 0. \tag{18}$$

We can multiply each term of (18) by $\frac{1}{\mu} \Delta(e^{-\frac{u}{2}}) - \frac{1}{2\mu} |\nabla u|^2 e^{-\frac{u}{2}}$ and integrate, which gives the following identities

$$\begin{aligned} - \int_{\mathbb{R}^2} \Delta u \Delta(e^{-u/2}) \frac{1}{\mu} dx &= \int_{\mathbb{R}^2} \Delta u \left(\Delta u - \frac{1}{2} |\nabla u|^2 \right) e^{-u/2} \frac{1}{2\mu} dx, \\ \int_{\mathbb{R}^2} \mu \frac{1}{\mu} \Delta(e^{-u/2}) dx &= 0, \\ \int_{\mathbb{R}^2} e^u \mu \left(\frac{1}{\mu} \Delta(e^{-\frac{u}{2}}) - \frac{1}{2\mu} |\nabla u|^2 e^{-\frac{u}{2}} \right) dx &= 0. \end{aligned}$$

Defining $\nu := e^{-u/2}/\mu = e^{-u/2-g}$ with $g := \log \mu$ and $d\nu := \nu dx$ we have

$$\mathcal{I}[u] = 2 \int_{\mathbb{R}^2} (\Delta u)^2 d\nu + \int_{\mathbb{R}^2} \Delta u |\nabla u|^2 d\nu - 16 \pi \lambda \int_{\mathbb{R}^2} |\nabla u|^2 e^{-u/2} dx = 0.$$

Let us introduce some notations, which are consistent with the ones on manifolds. Let us denote by $\text{Hu} = (\frac{\partial^2 u}{\partial x_i \partial x_j})_{i,j=1,2}$ the Hessian of u , $\text{Lu} = \text{Hu} - \frac{1}{2} \Delta u \text{I}_2$ is the trace free Hessian and $\text{M}[u] := \nabla u \otimes \nabla u - \frac{1}{2} |\nabla u|^2 \text{I}_2$, where $\nabla u \otimes \nabla u = (\frac{\partial u}{\partial x_i} \frac{\partial u}{\partial x_j})_{i,j=1,2}$. For the convenience of the reader, we split the computations in four steps.

1) Let us start with some preliminary computations. An integration by parts shows that

$$\begin{aligned} & 2 \int_{\mathbb{R}^2} \Delta u \nabla u \cdot \nabla g d\nu - \int_{\mathbb{R}^2} |\nabla u|^2 \nabla u \cdot \nabla g d\nu \\ &= -2 \int_{\mathbb{R}^2} \text{Hu} : (\nabla u \otimes \nabla g) d\nu - 2 \int_{\mathbb{R}^2} (\text{Hg} - \nabla g \otimes \nabla g) : (\nabla u \otimes \nabla u) d\nu \\ &= -2 \int_{\mathbb{R}^2} \text{Hu} : (\nabla u \otimes \nabla g) d\nu - 2 \int_{\mathbb{R}^2} \text{Hg} : (\nabla u \otimes \nabla u) d\nu + 2 \int_{\mathbb{R}^2} (\nabla u \cdot \nabla g)^2 d\nu. \end{aligned} \tag{19}$$

By expanding $\text{Lu} - \frac{1}{2} \text{M}[u]$, we also get that

$$\begin{aligned} & \int_{\mathbb{R}^2} \left(\text{Lu} - \frac{1}{2} \text{M}[u] \right) : (\nabla u \otimes \nabla g) d\nu \\ &= \int_{\mathbb{R}^2} \text{Hu} : (\nabla u \otimes \nabla g) d\nu - \frac{1}{2} \int_{\mathbb{R}^2} \Delta u \nabla u \cdot \nabla g d\nu - \frac{1}{4} \int_{\mathbb{R}^2} |\nabla u|^2 \nabla u \cdot \nabla g d\nu. \end{aligned} \tag{20}$$

Recalling the definition of $\nu = e^{-\frac{1}{2}u-g}$, we also find that

$$\begin{aligned} -\frac{1}{2} \int_{\mathbb{R}^2} |\nabla u|^2 \nabla u \cdot \nabla g d\nu &= \int_{\mathbb{R}^2} |\nabla u|^2 \nabla g e^{-g} \cdot \nabla (e^{-\frac{1}{2}u}) dx \\ &= -2 \int_{\mathbb{R}^2} \text{Hu} : (\nabla u \otimes \nabla g) d\nu + \int_{\mathbb{R}^2} |\nabla u|^2 (|\nabla g|^2 - \Delta g) d\nu. \end{aligned} \tag{21}$$

Equations (19), (20) and (21) allow us to eliminate

$$\int_{\mathbb{R}^2} \Delta u \nabla u \cdot \nabla g d\nu, \quad \int_{\mathbb{R}^2} |\nabla u|^2 \nabla u \cdot \nabla g d\nu \quad \text{and} \quad \int_{\mathbb{R}^2} \text{Hu} : (\nabla u \otimes \nabla g) d\nu$$

in terms of the other quantities:

$$\begin{aligned} \int_{\mathbb{R}^2} \Delta u \nabla u \cdot \nabla g d\nu &= \int_{\mathbb{R}^2} |\nabla u|^2 (|\nabla g|^2 - \Delta g) d\nu \\ &\quad - 2 \int_{\mathbb{R}^2} \left(\text{Lu} - \frac{1}{2} \text{M}[u] \right) : (\nabla u \otimes \nabla g) d\nu, \end{aligned} \tag{22}$$

$$\begin{aligned} \int_{\mathbb{R}^2} |\nabla u|^2 \nabla u \cdot \nabla g d\nu &= 4 \int_{\mathbb{R}^2} \text{Hg} : (\nabla u \otimes \nabla u) d\nu - 4 \int_{\mathbb{R}^2} (\nabla u \cdot \nabla g)^2 d\nu \\ &\quad - 8 \int_{\mathbb{R}^2} \left(\text{Lu} - \frac{1}{2} \text{M}[u] \right) : (\nabla u \otimes \nabla g) d\nu \\ &\quad + 6 \int_{\mathbb{R}^2} |\nabla u|^2 (|\nabla g|^2 - \Delta g) d\nu, \end{aligned} \tag{23}$$

$$\begin{aligned} \int_{\mathbb{R}^2} \mathbf{H}u : (\nabla u \otimes \nabla g) \, d\nu &= \int_{\mathbb{R}^2} \mathbf{H}g : (\nabla u \otimes \nabla u) \, d\nu - \int_{\mathbb{R}^2} (\nabla u \cdot \nabla g)^2 \, d\nu \\ &\quad - 2 \int_{\mathbb{R}^2} \left(\mathbf{L}u - \frac{1}{2} \mathbf{M}[u] \right) : (\nabla u \otimes \nabla g) \, d\nu \quad (24) \\ &\quad + 2 \int_{\mathbb{R}^2} |\nabla u|^2 (|\nabla g|^2 - \Delta g) \, d\nu. \end{aligned}$$

Moreover, the reader is invited to check that

$$\|\mathbf{L}u\|^2 = \|\mathbf{H}u\|^2 - \frac{1}{2} (\Delta u)^2 \quad (25)$$

and

$$\|\mathbf{M}[u]\|^2 = \frac{1}{2} |\nabla u|^4. \quad (26)$$

2) On the one hand, integrating the second term in the expression of $\mathcal{I}[u]$ by parts gives

$$\begin{aligned} &\int_{\mathbb{R}^2} \Delta u |\nabla u|^2 \, d\nu \\ &= - \int_{\mathbb{R}^2} \nabla u \cdot \nabla (|\nabla u|^2 \nu) \, dx \\ &= - \int_{\mathbb{R}^2} \nabla u \cdot \left(2 \mathbf{H}u \nabla u - \frac{1}{2} \nabla u |\nabla u|^2 - |\nabla u|^2 \nabla g \right) \, d\nu \\ &= - \int_{\mathbb{R}^2} \left(2 \mathbf{H}u : \nabla u \otimes \nabla u - \frac{1}{2} |\nabla u|^4 - |\nabla u|^2 (\nabla u \cdot \nabla g) \right) \, d\nu \\ &= - \int_{\mathbb{R}^2} \left(2 \mathbf{L}u : \nabla u \otimes \nabla u + \Delta u |\nabla u|^2 - \frac{1}{2} |\nabla u|^4 - |\nabla u|^2 (\nabla u \cdot \nabla g) \right) \, d\nu \\ &= - \int_{\mathbb{R}^2} \left(2 \mathbf{L}u : \mathbf{M}[u] + \Delta u |\nabla u|^2 - \frac{1}{2} |\nabla u|^4 - |\nabla u|^2 (\nabla u \cdot \nabla g) \right) \, d\nu, \end{aligned}$$

that is

$$\int_{\mathbb{R}^2} \Delta u |\nabla u|^2 \, d\nu = - \int_{\mathbb{R}^2} \left(\mathbf{L}u : \mathbf{M}[u] - \frac{1}{4} |\nabla u|^4 - \frac{1}{2} |\nabla u|^2 (\nabla u \cdot \nabla g) \right) \, d\nu.$$

According to (23) and (26), we obtain

$$\begin{aligned} \int_{\mathbb{R}^2} \Delta u |\nabla u|^2 \, d\nu &= - \int_{\mathbb{R}^2} \mathbf{L}u : \mathbf{M}[u] \, d\nu \\ &\quad + \frac{1}{2} \int_{\mathbb{R}^2} \|\mathbf{M}[u]\|^2 \, d\nu + 2 \int_{\mathbb{R}^2} \mathbf{H}g : (\nabla u \otimes \nabla u) \, d\nu \\ &\quad - 2 \int_{\mathbb{R}^2} (\nabla u \cdot \nabla g)^2 \, d\nu - 4 \int_{\mathbb{R}^2} \left(\mathbf{L}u - \frac{1}{2} \mathbf{M}[u] \right) : (\nabla u \otimes \nabla g) \, d\nu \\ &\quad + 3 \int_{\mathbb{R}^2} |\nabla u|^2 (|\nabla g|^2 - \Delta g) \, d\nu. \quad (27) \end{aligned}$$

3) On the other hand, integrating by parts twice yields

$$\int_{\mathbb{R}^2} \Delta |\nabla u|^2 \, d\nu = \int_{\mathbb{R}^2} |\nabla u|^2 \Delta (e^{-u/2}) \frac{dx}{\mu} + \int_{\mathbb{R}^2} |\nabla u|^2 (|\nabla g|^2 - \Delta g) \, d\nu$$

$$\begin{aligned}
 & + \int_{\mathbb{R}^2} |\nabla u|^2 (\nabla u \cdot \nabla g) \, d\nu, \\
 & = -\frac{1}{2} \int_{\mathbb{R}^2} \Delta u |\nabla u|^2 \, d\nu + \frac{1}{4} \int_{\mathbb{R}^2} |\nabla u|^4 \, d\nu \\
 & \quad + \int_{\mathbb{R}^2} |\nabla u|^2 (|\nabla g|^2 - \Delta g) \, d\nu + \int_{\mathbb{R}^2} |\nabla u|^2 (\nabla u \cdot \nabla g) \, d\nu. \tag{28}
 \end{aligned}$$

Integrating by parts again we have that

$$\int_{\mathbb{R}^2} \nabla \Delta u \cdot \nabla u \, d\nu = - \int_{\mathbb{R}^2} \Delta u \left(\Delta u - \frac{1}{2} |\nabla u|^2 - (\nabla u \cdot \nabla g) \right) \, d\nu,$$

which we can use along with the Bochner-Lichnerovicz-Weitzenböck formula on \mathbb{R}^2 (with Ricci tensor identically equal to 0),

$$\Delta |\nabla u|^2 = 2 \|Lu\|^2 + (\Delta u)^2 + 2 \nabla \Delta u \cdot \nabla u,$$

to get

$$\int_{\mathbb{R}^2} \Delta |\nabla u|^2 \, d\nu = \int_{\mathbb{R}^2} \left(2 \|Lu\|^2 - (\Delta u)^2 + \Delta u |\nabla u|^2 + 2 \Delta u (\nabla u \cdot \nabla g) \right) \, d\nu.$$

Combined with (28) this proves that

$$\begin{aligned}
 \int_{\mathbb{R}^2} (\Delta u)^2 \, d\nu & = 2 \int_{\mathbb{R}^2} \|Lu\|^2 \, d\nu + \frac{3}{2} \int_{\mathbb{R}^2} \Delta u |\nabla u|^2 \, d\nu - \frac{1}{4} \int_{\mathbb{R}^2} |\nabla u|^4 \, d\nu \\
 & - \int_{\mathbb{R}^2} |\nabla u|^2 (|\nabla g|^2 - \Delta g) \, d\nu + 2 \int_{\mathbb{R}^2} \Delta u (\nabla u \cdot \nabla g) \, d\nu - \int_{\mathbb{R}^2} |\nabla u|^2 (\nabla u \cdot \nabla g) \, d\nu.
 \end{aligned}$$

Using (22), (23), (26) and (27), we obtain

$$\begin{aligned}
 & \int_{\mathbb{R}^2} (\Delta u)^2 \, d\nu \\
 & = 2 \int_{\mathbb{R}^2} \|Lu\|^2 \, d\nu \\
 & \quad - \frac{3}{2} \int_{\mathbb{R}^2} Lu : M[u] \, d\nu + \frac{3}{4} \int_{\mathbb{R}^2} \|M[u]\|^2 \, d\nu + 3 \int_{\mathbb{R}^2} Hg : (\nabla u \otimes \nabla u) \, d\nu \\
 & \quad - 3 \int_{\mathbb{R}^2} (\nabla u \cdot \nabla g)^2 \, d\nu - 6 \int_{\mathbb{R}^2} \left(Lu - \frac{1}{2} M[u] \right) : (\nabla u \otimes \nabla g) \, d\nu \\
 & \quad + \frac{9}{2} \int_{\mathbb{R}^2} |\nabla u|^2 (|\nabla g|^2 - \Delta g) \, d\nu \\
 & \quad - \frac{1}{2} \int_{\mathbb{R}^2} \|M[u]\|^2 \, d\nu - \int_{\mathbb{R}^2} |\nabla u|^2 (|\nabla g|^2 - \Delta g) \, d\nu \\
 & \quad + 2 \int_{\mathbb{R}^2} |\nabla u|^2 (|\nabla g|^2 - \Delta g) \, d\nu - 4 \int_{\mathbb{R}^2} \left(Lu - \frac{1}{2} M[u] \right) : (\nabla u \otimes \nabla g) \, d\nu \\
 & \quad - 4 \int_{\mathbb{R}^2} Hg : (\nabla u \otimes \nabla u) \, d\nu + 4 \int_{\mathbb{R}^2} (\nabla u \cdot \nabla g)^2 \, d\nu \\
 & \quad + 8 \int_{\mathbb{R}^2} \left(Lu - \frac{1}{2} M[u] \right) : (\nabla u \otimes \nabla g) \, d\nu \\
 & \quad - 6 \int_{\mathbb{R}^2} |\nabla u|^2 (|\nabla g|^2 - \Delta g) \, d\nu.
 \end{aligned}$$

Collecting terms, we arrive at

$$\begin{aligned} \int_{\mathbb{R}^2} (\Delta u)^2 dv &= 2 \int_{\mathbb{R}^2} \|Lu\|^2 dv - \frac{3}{2} \int_{\mathbb{R}^2} Lu : M[u] dv + \frac{1}{4} \int_{\mathbb{R}^2} \|M[u]\|^2 dv \\ &\quad - 2 \int_{\mathbb{R}^2} \left(Lu - \frac{1}{2} M[u] \right) : (\nabla u \otimes \nabla g) dv \\ &\quad - \int_{\mathbb{R}^2} Hg : (\nabla u \otimes \nabla u) dv + \int_{\mathbb{R}^2} (\nabla u \cdot \nabla g)^2 dv \\ &\quad - \frac{1}{2} \int_{\mathbb{R}^2} |\nabla u|^2 (|\nabla g|^2 - \Delta g) dv. \quad (29) \end{aligned}$$

4) By reinjecting (27) and (29) in the expression of \mathcal{I} , we get that

$$\begin{aligned} 0 = \mathcal{I}[u] &= 4 \int_{\mathbb{R}^2} \|Lu\|^2 dv - 4 \int_{\mathbb{R}^2} Lu : M[u] dv + \int_{\mathbb{R}^2} \|M[u]\|^2 dv \\ &\quad - 8 \int_{\mathbb{R}^2} \left(Lu - \frac{1}{2} M[u] \right) : (\nabla u \otimes \nabla g) dv \\ &\quad + 2 \int_{\mathbb{R}^2} |\nabla u|^2 (|\nabla g|^2 - \Delta g) dv - 16\pi\lambda \int_{\mathbb{R}^2} |\nabla u|^2 e^{-u/2} dx. \end{aligned}$$

Since $\|\nabla u \otimes \nabla g\|^2 = |\nabla u|^2 |\nabla g|^2$, then we get for the corresponding trace free quantity

$$Nu := \nabla u \otimes \nabla g - \frac{1}{2} (\nabla u \cdot \nabla g) I_2$$

that

$$\|Nu\|^2 = |\nabla u|^2 |\nabla g|^2 - \frac{1}{2} (\nabla u \cdot \nabla g)^2$$

and hence obtain the identity

$$\begin{aligned} 0 &= 4 \int_{\mathbb{R}^2} \left\| Lu - \frac{1}{2} M[u] - Nu \right\|^2 dv \\ &\quad - 2 \int_{\mathbb{R}^2} [(\Delta g + |\nabla g|^2 - (\nabla g \cdot \omega)^2) e^{-g} + 8\pi\lambda] |\nabla u|^2 e^{-u/2} dx \quad (30) \end{aligned}$$

where $\omega := \nabla u / |\nabla u|$. If we assume that $\int_{\mathbb{R}^2} |\nabla u|^2 e^{-u/2} dx \neq 0$ and define

$$\Lambda := \frac{1}{8\pi} \frac{\int_{\mathbb{R}^2} [(\nabla u \cdot \nabla g)^2 - (\Delta g + |\nabla g|^2) |\nabla u|^2] e^{-u/2-g} dx}{\int_{\mathbb{R}^2} |\nabla u|^2 e^{-u/2} dx},$$

then we get a contradiction if $\lambda < \Lambda$. To keep maximal generality, one could even include the term $\int_{\mathbb{R}^2} \|Lu - \frac{1}{2} M[u] - Nu\|^2 dv$ in the definition of Λ , as it was done in the case of manifolds. However, this is a quite complicated criterion to verify since it involves the solution to (18) itself. Hence it makes sense to consider the simpler case where μ has radial symmetry. In that case it is also known from [32] that u is radially symmetric if μ is a monotone non-increasing function of $|x|$. Let

$$\Lambda_\star := -\frac{1}{8\pi} \inf_{x \in \mathbb{R}^2} (e^{-g} \Delta g) = \inf_{x \in \mathbb{R}^2} \frac{-\Delta \log \mu}{8\pi \mu}.$$

Theorem 8. *Assume that μ is a radially symmetric function. Then any radially symmetric solution to (18) is a constant if $\lambda < \Lambda_\star$ and the inequality (17) holds with $\lambda = \Lambda_\star$ if equality is achieved among radial functions.*

Example 1. The Euclidean Onofri inequality corresponds to

$$\mu(x) = \frac{1}{\pi(1+|x|^2)^2} \quad \forall x \in \mathbb{R}^2.$$

Since $-\Delta \log \mu = 8\pi\mu$, it is known that $\Lambda_\star = 1$ is the optimal constant. See [14] for further details. Let us notice that the analysis of the equation

$$Lu - \frac{1}{2}M[u] - Nu = 0$$

in the case $\Lambda_\star = 1$ provides a proof of the uniqueness of the radial solution to (18), which is alternative to the result of [11].

Example 2. It is straightforward to deduce a perturbation result from Theorem 8, that goes as follows. Let

$$\mu(x) = \frac{e^{-h(x)}}{Z(1+|x|^2)^2} \quad \forall x \in \mathbb{R}^2,$$

where h is a radial function and Z a normalization constant so that μ is a probability measure. We shall assume that h has a bounded variation and is such that $|x|^4 \Delta h$ is bounded from below. Then we have the estimate

$$\inf_{x \in \mathbb{R}^2} \frac{-\Delta \log \mu}{8\pi\mu} \geq e^{-\text{Var}(h)} \left[1 + \frac{1}{8} \inf_{x \in \mathbb{R}^2} (1+|x|^2)^2 \Delta h \right].$$

Example 3. The subcritical Onofri inequality has been studied in [12]. It plays an important role for the study of the subcritical Keller-Segel model and its asymptotics for large times, and goes as follows. Let $\mu = n/M$ where n is given as the unique (up to constants) radial solution to

$$-\Delta c = n = M \frac{e^{c-\frac{1}{2}|x|^2}}{\int_{\mathbb{R}^2} e^{c-\frac{1}{2}|x|^2} dx}$$

and the mass M is taken in the interval $M \in (0, 8\pi)$. According to [32], n is radially symmetric as a consequence of the method of moving planes. It is straightforward to check that

$$\inf_{x \in \mathbb{R}^2} \frac{-\Delta \log \mu}{8\pi\mu} = \frac{M}{8\pi} + \inf_{x \in \mathbb{R}^2} \frac{1}{4\pi\mu}$$

but the symmetry of the solution to (17) is not true for $\lambda > \frac{M}{8\pi}$ and $\frac{M}{8\pi}$ turns out to be the value of the optimal constant in (17). See [13, 12] for further details.

Example 4. The parabolic-parabolic Keller-Segel model has global in time solutions with mass larger than 8π for some values of its parameters, according to [10]. The stationary solutions in self-similar variables can be written as

$$-\Delta c = \varepsilon x \cdot \nabla c + n \quad \text{with} \quad n = M \frac{e^{c-\frac{1}{2}|x|^2}}{\int_{\mathbb{R}^2} e^{c-\frac{1}{2}|x|^2} dx}$$

(where $\varepsilon > 0$ is a given parameter) and have been shown to be radially symmetric in [39]. To prove that a weighted Onofri inequality holds with $\mu = n/M$, it is

therefore sufficient to establish the range of $\lambda \in (0, \Lambda_*)$ such that the minimizer of $u \mapsto \int_{\mathbb{R}^2} |\nabla u|^2 dx - 16\pi\lambda [\log(\int_{\mathbb{R}^2} e^u d\mu) - \int_{\mathbb{R}^2} u d\mu]$ is radially symmetric, where

$$\inf_{x \in \mathbb{R}^2} \frac{-\Delta \log \mu}{8\pi\mu} = \frac{M}{8\pi} + \inf_{x \in \mathbb{R}^2} \frac{\varepsilon x \cdot \nabla c + 2}{8\pi\mu}.$$

Such symmetry breaking issues are however known to be difficult: see for instance [23] for a discussion of a related problem.

Acknowledgments. J.D. and G.J. have been supported by the projects *STAB* and *Kibord* of the French National Research Agency (ANR). J.D. and M.J.E. have been supported by the project *NoNAP* of the French National Research Agency (ANR). The authors thank M. Loss and M. Kowalczyk for many stimulating discussions.

REFERENCES

- [1] T. Aubin, Meilleures constantes dans le théorème d'inclusion de Sobolev et un théorème de Fredholm non linéaire pour la transformation conforme de la courbure scalaire, *J. Funct. Anal.*, **32** (1979), 148–174.
- [2] T. Aubin, *Nonlinear Analysis on Manifolds. Monge-Ampère Equations*, vol. 252 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], Springer-Verlag, New York, 1982.
- [3] D. Bakry and M. Ledoux, Sobolev inequalities and Myers's diameter theorem for an abstract Markov generator, *Duke Math. J.*, **85** (1996), 253–270.
- [4] D. Bakry, I. Gentil and M. Ledoux, *Analysis and Geometry of Markov Diffusion Operators*, vol. 348 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], Springer, Cham, 2014.
- [5] W. Beckner, Sharp Sobolev inequalities on the sphere and the Moser-Trudinger inequality, *Ann. of Math. (2)*, **138** (1993), 213–242.
- [6] A. Bentaléb, Inégalité de Sobolev pour l'opérateur ultrasphérique, *C. R. Acad. Sci. Paris Sér. I Math.*, **317** (1993), 187–190.
- [7] M. Berger, P. Gauduchon and E. Mazet, *Le Spectre d'une Variété Riemannienne*, Lecture Notes in Mathematics, Vol. 194, Springer-Verlag, Berlin, 1971.
- [8] R. J. Berman and B. Berndtsson, Moser-Trudinger type inequalities for complex Monge-Ampère operators and Aubin's "hypothèse fondamentale", ArXiv e-prints.
- [9] M.-F. Bidaut-Véron and L. Véron, Nonlinear elliptic equations on compact Riemannian manifolds and asymptotics of Emden equations, *Invent. Math.*, **106** (1991), 489–539.
- [10] P. Biler, L. Corrias and J. Dolbeault, Large mass self-similar solutions of the parabolic-parabolic Keller–Segel model of chemotaxis, *Journal of Mathematical Biology*, **63** (2011), 1–32.
- [11] P. Biler, G. Karch, P. Laurençot and T. Nadzieja, The 8π -problem for radially symmetric solutions of a chemotaxis model in the plane, *Math. Methods Appl. Sci.*, **29** (2006), 1563–1583.
- [12] J. Campos and J. Dolbeault, A functional framework for the Keller-Segel system: Logarithmic Hardy-Littlewood-Sobolev and related spectral gap inequalities, *C. R. Math. Acad. Sci. Paris*, **350** (2012), 949–954.
- [13] J. F. Campos and J. Dolbeault, Asymptotic estimates for the parabolic-elliptic keller-segel model in the plane, *Comm. Partial Differential Equations*, **39** (2014), 806–841.
- [14] E. A. Carlen and M. Loss, Competing symmetries, the logarithmic HLS inequality and Onofri's inequality on \mathbb{S}^n , *Geom. Funct. Anal.*, **2** (1992), 90–104.
- [15] S.-Y. A. Chang, Extremal functions in a sharp form of Sobolev inequality, in *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986)*, Amer. Math. Soc., Providence, RI, 1987, 715–723.
- [16] S.-Y. A. Chang and P. C. Yang, Conformal deformation of metrics on \mathbb{S}^2 , *J. Differential Geom.*, **27** (1988), 259–296.
- [17] S.-Y. A. Chang, *Non-linear Elliptic Equations in Conformal Geometry*, Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2004.
- [18] S.-Y. A. Chang and P. C. Yang, Prescribing Gaussian curvature on \mathbb{S}^2 , *Acta Math.*, **159** (1987), 215–259.

- [19] S.-Y. A. Chang and P. C. Yang, The inequality of Moser and Trudinger and applications to conformal geometry, *Comm. Pure Appl. Math.*, **56** (2003), 1135–1150, Dedicated to the memory of Jürgen K. Moser.
- [20] W. X. Chen and C. Li, Classification of solutions of some nonlinear elliptic equations, *Duke Math. J.*, **63** (1991), 615–622.
- [21] P. Cherrier, Une inégalité de Sobolev sur les variétés riemanniennes, *Bull. Sci. Math. (2)*, **103** (1979), 353–374.
- [22] J. Demange, Improved Gagliardo-Nirenberg-Sobolev inequalities on manifolds with positive curvature, *J. Funct. Anal.*, **254** (2008), 593–611.
- [23] J. Dolbeault and M. J. Esteban, Branches of non-symmetric critical points and symmetry breaking in nonlinear elliptic partial differential equations, *Nonlinearity*, **27** (2014), 435–465.
- [24] J. Dolbeault, M. J. Esteban and G. Jankowiak, The Moser-Trudinger-Onofri inequality, *Chinese Annals of Math. B*, **36** (2015), 777–802.
- [25] J. Dolbeault, M. J. Esteban, M. Kowalczyk and M. Loss, Improved interpolation inequalities on the sphere, *Discrete and Continuous Dynamical Systems Series S (DCDS-S)*, **7** (2014), 695–724.
- [26] J. Dolbeault, M. J. Esteban and M. Loss, Nonlinear flows and rigidity results on compact manifolds, *Journal of Functional Analysis*, **267** (2014), 1338–1363.
- [27] Z. Faget, Best constants in the exceptional case of Sobolev inequalities, *Math. Z.*, **252** (2006), 133–146.
- [28] Z. Faget, Optimal constants in the exceptional case of Sobolev inequalities on Riemannian manifolds, *Trans. Amer. Math. Soc.*, **360** (2008), 2303–2325.
- [29] É. Fontenas, Sur les constantes de Sobolev des variétés riemanniennes compactes et les fonctions extrémales des sphères, *Bull. Sci. Math.*, **121** (1997), 71–96.
- [30] A. Ghigi, On the Moser-Onofri and Prékopa-Leindler inequalities, *Collect. Math.*, **56** (2005), 143–156.
- [31] N. Ghoussoub and A. Moradifam, *Functional Inequalities: New Perspectives and New Applications*, vol. 187 of Mathematical Surveys and Monographs, American Mathematical Society, Providence, RI, 2013.
- [32] B. Gidas, W. M. Ni and L. Nirenberg, Symmetry of positive solutions of nonlinear elliptic equations in \mathbb{R}^n , in *Mathematical analysis and applications, Part A*, vol. 7 of Adv. in Math. Suppl. Stud., Academic Press, New York-London, 1981, 369–402.
- [33] E. Hebey, *Nonlinear Analysis on Manifolds: Sobolev Spaces and Inequalities*, vol. 5 of Courant Lecture Notes in Mathematics, New York University Courant Institute of Mathematical Sciences, New York, 1999.
- [34] C. W. Hong, A best constant and the Gaussian curvature, *Proc. Amer. Math. Soc.*, **97** (1986), 737–747.
- [35] Y. Y. Li, Harnack type inequality: The method of moving planes, *Comm. Math. Phys.*, **200** (1999), 421–444.
- [36] A. Lichnerowicz, *Géométrie des Groupes de Transformations*, Travaux et Recherches Mathématiques, III. Dunod, Paris, 1958.
- [37] J. R. Licois and L. Véron, Un théorème d’annulation pour des équations elliptiques non linéaires sur des variétés riemanniennes compactes, *C. R. Acad. Sci. Paris Sér. I Math.*, **320** (1995), 1337–1342.
- [38] J. Moser, A sharp form of an inequality by N. Trudinger, *Indiana Univ. Math. J.*, **20** (1970/71), 1077–1092.
- [39] Y. Naito, T. Suzuki and K. Yoshida, Self-similar solutions to a parabolic system modeling chemotaxis, *J. Differential Equations*, **184** (2002), 386–421.
- [40] M. Nolasco and G. Tarantello, On a sharp Sobolev-type inequality on two-dimensional compact manifolds, *Arch. Ration. Mech. Anal.*, **145** (1998), 161–195.
- [41] E. Onofri, On the positivity of the effective action in a theory of random surfaces, *Comm. Math. Phys.*, **86** (1982), 321–326.
- [42] B. Osgood, R. Phillips and P. Sarnak, Extremals of determinants of Laplacians, *J. Funct. Anal.*, **80** (1988), 148–211.
- [43] D. H. Phong, J. Song, J. Sturm and B. Weinkove, The Moser-Trudinger inequality on Kähler-Einstein manifolds, *American Journal of Mathematics*, **130** (2008), 1067–1085.
- [44] Y. A. Rubinstein, On energy functionals, Kähler-Einstein metrics, and the Moser-Trudinger-Onofri neighborhood, *J. Funct. Anal.*, **255** (2008), 2641–2660.

- [45] Y. A. Rubinstein, Some discretizations of geometric evolution equations and the Ricci iteration on the space of Kähler metrics, *Adv. Math.*, **218** (2008), 1526–1565.
- [46] G. Tarantello, *Seldual Gauge Field Vortices: An Analytical Approach*, Progress in Nonlinear Differential Equations and their Applications, 72, Birkhäuser Boston, Inc., Boston, MA, 2008.
- [47] G. Tian and X. Zhu, A nonlinear inequality of Moser-Trudinger type, *Calculus of Variations and Partial Differential Equations*, **10** (2000), 349–354.
- [48] N. S. Trudinger, On imbeddings into Orlicz spaces and some applications, *J. Math. Mech.*, **17** (1967), 473–483.

Received November 2015; revised January 2017.

E-mail address: `dolbeaul@ceremade.dauphine.fr`

E-mail address: `esteban@ceremade.dauphine.fr`

E-mail address: `gaspard.jankowiak@math.cnrs.fr`